arXiv:1304.4545v1 [astro-ph.SR] 16 Apr 2013

Correlation Study Of Some Solar Activity Indices In The Cycles 21 - 23

E.A. Bruevich [a] , V. V. Bruevich [b]
[a]Sternberg Astronomical Institute, Moscow State University, Russia
[b]Sternberg Astronomical Institute, Moscow State University, Russia
E-mail: [a]red-field@yandex.ru, [b]brouev@sai.msu.ru

Abstract.

The correlation coefficients of the linear regression of six solar indices versus $F_{10,7}$ were analyzed in solar cycles 21, 22 and 23. We also analyzed the interconnection between these indices and $F_{10,7}$ with help of the approximation by the polynomials of second order. The indices we've studied in this paper are: Wolf numbers - W, 530,3 nm coronal line flux - F_{530}, the total solar irradiance - TSI, Mg II UV-index 280 nm core-to-wing ratio, Flare Index - FI and Counts of flares. In the most cases the regressions of these solar indices versus $F_{10,7}$ are close to linear except the moments of time near to the minimums and maximums of 11-year activity. For the linear regressions we found that the values of correlation coefficients $K_{corr}(t)$ for the indices versus $F_{10,7}$ and W show the cyclic variations with periods approximately equal to the to half length of 11-year cycle - 5,5 years approximately.

KEY WORDS: Solar cycle, Observations, Solar activity indices

1 Introduction

Magnetic activity of the Sun is called the complex of electromagnetic and hydrodynamic processes in the solar atmosphere. The analysis of active regions (plages and spots in the photosphere, flocculae in the chromosphere and prominences in the corona of the Sun) is required to study the magnetic field of the Sun and the physics of magnetic activity. This task is of fundamental importance for astrophysics of the Sun and the stars. Its applied meaning is connected with the influence of solar active processes on the Earth's magnetic field.

We have studied monthly averaged values of six global solar activity indices in magnetic activity cycles 21, 22 and 23. Most of these observed data we used in our paper were published in Solar-Geophysical Data Reports (2009) and National Geophysical Data Center Solar and Terrestrial Physics (2012).

All the indices studied in this paper are very important not only for analysis of solar radiation formed on the different altitudes of solar atmosphere, but for solar-terrestrial relationships as the key factors of the solar radiation influence (EUV/UV solar radiation is the most important) on the different layers of terrestrial atmosphere also.

It's known that the various solar activity parameters correlate quite well with the most popular solar index sunspot numbers W and with each others over long time scales. Floyd et al. (2005) showed that the mutual relation between sunspot numbers and three solar UV/EUV indices, the $F_{10,7}$ flux, the MgII core-to-wing ratio, which is the important chromospheric UV index (Viereck et al. (2001), Viereck et al. (2004)), remained stable for 25 years until 2000. At the end of 2001 these mutual relations dramatically changed due to a large enhancement which took place after actual sunspot maximum of the cycle 23 and the subsequent relative quietness intermediate called the Gnevyshev gap.

In our issue for all the indices we used the monthly averages values. Such averages allowed us to take into consideration the fact that the major modulation of solar indexes contains a periodicity of about 27-28 days (corresponding to the mean solar rotation period). So we reduced the influence of the rotational modulation of the data sets.

Vitinsky et al. (1986) analyzed solar cycles 18 - 20 and pointed out that correlation for spot numbers versus radio flux $F_{10,7}$ does not show the close linear connection during all the activity cycle. Also it was emphasized the importance of statistical study in our solar activity processes understanding. To achieve the best agreement in approximation of spot numbers values by $F_{10,7}$ observations Vitinsky et al. (1986) proposed to approximate the dependence W - $F_{10,7}$ by two linear regressions: the first one - for the low solar activity (where $F_{10,7}$ less than 150) and the second one - for the high activity ($F_{10,7}$ more than 150).

In this paper we found out that the linear correlation was violated not only for maximums of solar activity cycles but for minimums of the cycles too. Our analysis of the interconnection between these indices and $F_{10,7}$ with help of the approximation by the polynomials of second order confirmed this fact.

We also analyzed a three-year determined correlation coefficients $K_{corr}(t)$ for six solar activity indices versus $F_{10,7}$.

2 Global activity indices

Then we have to say a few words about solar indices studied in this paper.

At Figure 1 we show the activity indices which are normalized to their values averaged over the analyzed time interval. We can see that the relative variation of the index in a solar cycle is about 2-3 times. However, the



Figure 1: The time series of monthly average values of W, $F_{10,7}$, MgII core-to-wing ratio, $F_{530,3}$, Flare Index and Counts of flares. The upper index N indicates that solar activity indices are normalized to their values averaged over the analyzed time interval.

magnitude of MgII - index (as well as TSI , not shown in this figure) change very little - about shares of percent.

The solar radio microwave flux at wavelengths 10,7 cm $F_{10,7}$ has the long running series of observations started in 1947 in Ottawa, Canada and maintained to this day at Penticton site in British Columbia. This radio emission comes from high part of the chromosphere and low part of the corona. $F_{10,7}$ radio flux has two different sources: thermal bremsstrahlung (due to electrons radiating when changing direction by being deflected by other charged participles) and gyro-radiation (due to electrons radiating when changing direction by gyrating around magnetic fields lines). These mechanisms give rise to enhanced radiation when the temperature, density and magnetic fields are enhanced. So $F_{10,7}$ is a good measure of general solar activity. $F_{10,7}$ data are



Figure 2: The time series of the average annual W from 1700 for 2009, and its wavelet image (we use the Morley wavelet transfer function).

available at http://radbelts.gsfc.nasa.gov/RB_model_model_int/Psi_database.html.

The MgII 280 nm is important solar activity indicator of radiation, derived from daily solar observations of the core-to-wing ratio of the MgII doublet at 279,9 nm provides a good measure of the solar UV variability and can be used as a reliable proxy to model extreme UV (EUV) variability during the solar cycle Scupin et al. (2005). The MgII observation data were obtained from several satellite's (NOAA, ENVISAT) instruments. NOAA started in 1978 (during the 21^{st}, 22^{nd} and the first part of the 23^{rd} solar activity cycles), ENVISAT was launched on 2002 (last part of the 23^{th} solar activity cycle). Comparison of the NOAA and ENVISAT MgII index observation data shows that both the MgII indexes agree to within about 0,5%. We used both the NOAA and ENVISAT MgII index observed data from Puga and Viereck (2004), Scupin et al. (2005). Viereck et al. (2001) showed an extremely good fit between the 30,4 nm emission (the main component of EUV-emission) and the NOAA MgII index. Viereck et al. (2001), Floyd et al. (2005) have noted the linear relationship between the MgII index and total solar irradiance.

Solar irradiance is the total amount of solar energy at a given wavelength received at the top of the earth's atmosphere per unit time. When integrated over all wavelengths, this quantity is called the total solar irradiance (TSI) previously known as the solar constant. Regular monitoring of TSI has been carried out since 1978. From 1985 the total solar irradiance was observed by Earth Radiation Budget Satellite (EBRS). We use the TSI data set from NGDC web site available at http://www.ngdc.noaa.gov/stp and combined observational data from National Geophysical Data Center Solar and Terrestrial Physics (2012). The importance of UV/EUV influence to TSI variability (Active Sun/Quiet Sun) was pointed by Krivova and Solanki (2008). There were indicated that up to 63,3 % of TSI variability is produced at wavelengths below 400 nm. Towards activity maxima the number of sunspots grows dramatically. But on average the Sun brightens about 0,1%. This is due to the increase amounts of bright features, faculae and network elements on the solar surface. The total area of the solar surface covered by such features rises more strongly as the cycle progresses than area of dark sunspots. Some physics-based models have been developed with using the combined proxies describing sunspot darkening (sunspot number or areas) and facular brightening (facular areas, CaII or MgII indices), see Frontenla et al. (2004), Krivova et al. (2003).

We also analyzed two activity indices which describe rapid processes on the Sun - Flare Index (FI) and monthly Counts of grouped solar flares (Monthly flare count). According to Solar-Geophysical Data Reports (2009) the term 'grouped' means observations of the same event by different sites were lumped together and counted as one. Kleczek (1952) defined the value $FI = it$ to quantify the daily flare activity over 24 hours per day. He assumed that relationship roughly gave the total energy emitted by the flare and named it flare index (FI). In this relation i represents the intensity scale of importance of the flare and and t the duration of the flare in minutes. In this issue we used the monthly averaged FI values.

The Wolf numbers is a very popular, widely used solar activity index: the series of Wolf numbers observations continue more than two hundred years. At Figure 2 we illustrated with help of wavelet - analysis the fact that the long time series of observations give us the very useful information for study of the problem of solar flux cyclicity on long time scales. The result of wavelet - analysis (Morley wavelet) of series of observations of average annual W is presented in form of many of isolines. For the isoline of the value of the wavelet-coefficients are of the same. The maximum values of isolines specify the maximum values of wavelet-coefficients, which corresponds to the most likely value of the period of the cycle. There are three well-defined cycles of activity: - the main cycle of activity is approximately equal to a 10 - 11 years, 40-50- year cyclicity and 100 to 120-year (ancient) cyclicity.

3 Activity indices in the cycles 21, 22 and in unusual cycle 23

The recent solar cycle 23 was the outstanding cycle for authentic observed data from 1849 year. It lasted 12,7 years and was the longest one for two hundred years of direct solar observations. This cycle is the second component in the 22-year Hale magnetic activity cycle but the 23^{rd} cycle was the first case of modern direct observations (from 1849 to 2008 years) when Gnevyshev-Ol's rule was violated: activity indices in cycle 23 had their maximum values less then the values in cycle 22 (but according to Gnevyshev-Ol's rule the cycle 23 must dominate), see Figure 2.

Ishkov (2009) pointed that in this unusual cycle 23 the monthly averages values for Wolf numbers during 8 months exceeded 113 and most of sunspot groups were less in size, their magnetic fields were less composite and characterized by the greater lifetime near 2^{nd} maximum in comparison with values near the 1^{st} maximum.The Wolf numbers reaches its first maximum 3.9 yrs after the beginning. After the first maximum, the index decrease by 14% (of that maximum). The two maxima of this index have the same amplitude

In the cycle 23 the $F_{10,7}$ radio flux and the TSI have the lowest values from 2007 to 2009 (the beginning of the cycle 24) all over of these indices observation period. The $F_{10,7}$ radio flux index shows the second maximum is 8.4% stronger than the first one. The TSI (from Earth Radiation Budget Satellite) maxima are fainter than those of the other indices because the solar irradiance variation of the 23rd solar cycle is just 0.14%. This value seems very small but is normal.

The Flare index - counts of flares has a higher first maximum. This shows that the flares can be generated during the first maximum, and it seems that the generation is decreasing towards the end of the cycle.

Figures 1 demonstrates that for all activity indices in the 23^{rd} solar activity cycle we can see two maximums separated one from another on 1,5 year approximately. We see the similar double-peak structure in cycle 22 but for the cycle 21 the double-peak structure is not evident. We see that there are displacements in both maximum occurrence time of all these indices in the 23^{rd} solar cycle.



Figure 3: Correlation between monthly averages for solar indices versus $F_{10,7}$ radio flux in the cycles 21 - 23. (a) $F_{530,3}$ coronal line flux, (b) MgII core-to-wing ratio, (c) TSI and (d) Counts of flares.

We assume that the probable reason of such double-peak structures is a modulation of the 11-year fluxes variations by both of the quasi-biennial and 5,5 year cyclicity. The different time of 1^{st} and 2^{nd} maximums appearance may be caused by the difference in fluxes formation conditions (for our indices) at different atmosphere's altitudes of the Sun.

Figures 1 also shows that for all solar indices in the cycle 23 the relative depth of the cavity between two maximums is about $10 - 15\%$.

When studied solar activity indices in 21, 22 and 23 solar activity cycles we separated out rise phases, cycle's maximum phase, cycle's minimum phase and decline phase. In case of linear regression we've found that the maximum values of correlation coefficients K_{corr} reached for the rise and decline phases of the cycles. According to our calculations the highest values of correlation coefficients K_{corr} we see in connection between W and $F_{10,7}$. Correlation coefficients K_{corr} for linear regression for TSI versus $F_{10,7}$ are the minimal of all correlation coefficients determined here. Figure 3 also illustrates the high level of interconnection between 4 solar activity indices versus $F_{10,7}$. Also we see that coefficients of linear regression (slope and intercept) differ among themselves for the activity cycles 21 - 23.

The cyclic variation of fluxes in different spectral ranges and lines at the 11-year time scale are widely spread phenomenon for F, G and K stars (not only for the Sun), see Bruevich & Kononovich (2011). The chromospheric activity indices (radiative fluxes at the centers of the H and K emission lines of Ca II - $396,8$ and $393,4$ nm respectively) for solar-type stars were studied during HK project by Baliunas (1995) at Mount Wilson observational program during 45 years, from 1965 to the present time. Authors of the HK project supposed that all the solar-type stars with well determined cyclic activity about 25% of the time remain in the Maunder minimum conditions. Some scientists proposed that the solar activity in future cycle 24 will be very low and similar to activity during the Maunder minimum period. Unlike this mention Chumak & Matveychuk (2010) predicted the main parameters of the new 24^{th} cycle closed to the usual activity cycle's parameters not similar to the cycle's characteristics during the Maunder minimum. At Figure 2 for yearly W we also see (if we continue the imaginary line, enveloped the maximums) the influence of the century cyclicity to the 24^{th} cycle's W maximum values.

4 Changed relation between activity indices and $F_{10,7}$ in the cycles 21-23

We analyzed the interconnection between activity indices W, $F_{530,3}$, MgII core to wing ratio, Flare Index, Counts of flares and TSI versus $F_{10,7}$ for the 21^{st}, 22^{nd} and 23^{th} solar cycles. At Figure 4 we presented the interconnection between solar indices and radio flux $F_{10,7}$ in the cycles 21 - 23.

The linear model corresponds to the linear regression equation:

$$F_{ind} = A_{ind} + B_{ind} \cdot F_{10,7} \, . \quad (1)$$

were F_{ind} is the activity index flux,
A_{ind} is the intercept of a linear regression,
B_{ind} is the slope of a linear regression.

Table 1: Solar activity indices versus $F_{10,7}$. Coefficients of linear regressions: A, B and their standard errors.

Activity indices versus $F_{10,7}$	A_{ind}	B_{ind}	Error σ_A	Error σ_B
W	-62,28	1,03	1,57	0,011
$F_{530,3}$	-2,93	0,084	0,27	0,002
MgII	0,258	0,00013	0,00026	0,00000018
Flare Index	-7,37	0,106	0,51	0,0036
Counts of flares	-269,03	4,31	20,14	0,146
TSI	1365,07	0,0066	0,044	0,0003

In Table 1 we present the coefficients of the linear regressions and their standard errors σ of intercept and slope values.

Ishkov (2009) pointed that there was very high level of flared activity in cycle 21 and very low level of flared activity in cycle 23. We also confirm this fact (see Figure 3d) that the difference of values $A_{Countsofflares}$ and $B_{Countsofflares}$ determined in 23^{rd} cycle from similar values, determined for cycles 21 and 22, is more significant among all other indices. The flared activity in the 23^{rd} cycle almost twice weaker in comparison to 21^{st} cycle.

Table 2: Solar activity indices versus $F_{10,7}$. Coefficients of polynomial regressions: A, $B1$, $B2$ and their standard errors.

Activity indices versus $F_{10,7}$	A	B1	B2	error σ_A	error σ_{B1}	error σ_{B2}
W	-87,26	1,45	-0,0015	4,86	0,078	0,0003
$F_{530,3}$	-7,38	0,158	-0,00026	0,84	0,013	0,000048
MgII	0,25	0,0002	-0,0000003	0,0007	0,00001	0,00000004
Flare Index	-4,36	0,057	0,00017	1,58	0,024	0,00007
Counts of flares	-361,4	5,83	-0,005	64,01	1,01	0,0035
TSI	1364,4	0,017	-0,000037	0,13	0,002	0,0000007

The polynomial model corresponds to the following equation of a second order polynomial:

$$F_{ind} = A_{ind} + B1_{ind} \cdot F_{10,7} + B2_{ind} \cdot F_{10,7}^2 \,. \qquad (2)$$

were F_{ind} is the activity index flux,

A_{ind} is the intercept of a polynomial regression,

$B1_{ind}$ and $B2_{ind}$ are the coefficients of a polynomial regression.

In Table 2 we present the coefficients of polynomial regressions: A, $B1$, $B2$ and their standard errors σ of intercept and slope values.

We have to point out that close interconnection between radiation fluxes characterized the energy release from different atmosphere's layers is the widespread phenomenon among the stars of late-type spectral classes. Bruevich & Alekseev (2007) confirmed that there exists the close interconnection between photospheric and coronal fluxes variations for solar-type stars of F, G, K and M spectral classes with widely varying activity of their atmospheres. It was shown that the summary areas of spots and values of X-ray fluxes increase gradually from the sun and HK project stars with the low spotted discs to the highly spotted K and M-stars for which Alekseev & Gershberg (1996) constructed the zonal model of the spots distributed at the star's disks .

We've calculated values K_{corr} of linear regression for solar activity indices versus $F_{10,7}$ for cycles 21,22 and 23. The values $K_{corr}(t)$ were determined for each moment of time t from K_{corr} calculation during the time interval $t - 1,5yr \ < \Delta T < \ t + 1,5yr$.

Figure 5 demonstrates the results of our correlation calculations of these solar activity indices versus $F_{10,7}$ (K_{corr} variations during the cycles 21 - 23). We can see that all the K_{corr} values have the maximum amounts at rise and at decline phases. It's easy to estimate the value of period of K_{corr} cyclic variations as 5,5 years approximately. We assumed that this new cyclicity (characterized with period's value equal to half length of 11-year cycle) is important for the successful forecasts of the solar activity indices fluxes.

The cyclic behavior of K_{corr} can be explained by following assumption: we imagine that some activity index flux depends on time t by the expression:

$$F_{ind}(t) = F_{ind}^{background}(t) + \Delta F_{ind}^{AR}(t) \,. \qquad (3)$$

were $F_{ind}^{background}(t)$ is the background flux rising continuously with increasing of solar activity and $\Delta F_{ind}^{AR}(t)$ is the additional flux to the overall flux from the active regions.

The previous correlation study allows us to consider that $F_{ind}^{background}(t)$ and $\Delta F_{ind}^{AR}(t)$ are the linear functions of the background and activity regions levels of solar activity. In our case we choose the radio flux $F_{10,7}$ as the best basal indicator of solar activity levels:

$$F_{ind}^{background}(t) = a_1 + b_1 \cdot F_{10,7}^{background}(t)\,. \tag{4}$$

$$\Delta F_{ind}^{AR}(t) = a_2 + b_2 \cdot \Delta F_{10,7}^{AR}(t)\,. \tag{5}$$

The coefficients a_1 and b_1 vary from a_2 and b_2 in different power for our different activity indices. For Wolf numbers this difference is small, but for 0,1 - 0,8 nm background and count of flares the difference between a_1, b_1 and a_2, b_2 is more significant than for W and TSI.

During the rise and decline cycle's phases the dependence $F_{ind}(t)$ versus $F_{10,7}(t)$ is approximately linear because coefficients a and b from Table 2 (which described all the cycle) are close to a_1 and b_1 (see Figure 3 and Figure 4) and relative addition flux from active regions $\Delta F_{ind}^{AR}(t)$ is neglect with respect to $F_{ind}^{background}(t)$. So additional flux from active regions cannot destroy a balance in the close linear correlation between $F_{ind}(t)$ and $F_{10,7}(t)$ and respective values of K_{corr} reach their maximum from all the cycle.

During the minimum of activity cycle both values $F_{ind}^{background}(t)$ and $\Delta F_{ind}^{AR}(t)$ are small, but additional flux from active regions is not neglect in relation to background flux that has the minimum values from all the activity cycle. Therefore values a and b from Table 2 cannot describe the linear regression to a considerable degree in cycle's minimum and values of K_{corr} reach their minimum values in the cycle.

During the maximum of activity cycle $\Delta F_{ind}^{AR}(t)$ often exceeds $F_{ind}^{background}(t)$ so disbalance in linear regression between activity indices increases and values of K_{corr} reach their minimum values in the activity cycle too.

5 Conclusions

For a long time the scientists were interested in the simulation of processes in the earth's ionosphere and upper atmosphere. It's known that the solar radiance at 30,4 nm is very significant for determination of the Earth high thermosphere levels heating. Lukyanova & Mursula (2011) showed that the for solar 30,4 nm radiance fluxes forecasts (very important for Earth thermosphere's heating predictions) there were more prefer to use MgII 280 nm observed data unlike usual $F_{10,7}$ and Wolf number observations.

The close interconnection between activity indices make possible new capabilities in the solar activity indices forecasts. For the successful forecasts of maximum values and other parameters of future activity cycles it has been required to take into account the century component. At Figure 2 we see the smoothed curved described the maximums of cycles which shows the century cyclicity dependence on Wolf numbers cycle's maximums.

In this paper we found out the cyclic behavior of a calculated during three-year interval values of correlation coefficients $K_{corr}(t)$ of linear regression for W, total solar irradiance, MgII 280 nm and Counts of flares versus $F_{10,7}$ during solar activity cycles 21,22 and 23 (see Figure 5). We showed that $K_{corr}(t)$ have the maximum values at the rise and decline phases - the linear connection between indices is more strong in these cases. It means that the forecasts of solar indices, based on $F_{10,7}$ observations in our case, will be more successful during the rise and decline cycle's phases.

We also determined that a calculated during three-year interval values of correlation coefficients $K_{corr}(t)$ are characterized by cyclic variations with the period that is equal to half length of period at 11 - year time scale (5,5 years about). Our study of linear regression between solar indices and $F_{10,7}$ confirms the fact that at minimum and at maximum cycle's phases the non-linear state of interconnection between solar activity indices (characterized the energy release from different layers of solar atmosphere) increases.

Acknowledgements The authors thank the RFBR grant 12-02-00884 for support of the work.

References

Alekseev, I.Yu. & Gershberg, R.E. 1996, On spotting of red dwarf stars: direct and inverse problem of the construction of zonal model. Astronomy Report. V. 73 pp. 589–597.

Baliunas, S.L., & Donahue, R.A., & Soon, W.H. 1995, Chromospheric variations in main-sequence stars. Astrophysical Journal, V. 438, pp. 269–280.

Bruevich, E.A., & Alekseev I.Yu. 2007, Spotting in stars with a low level of activity, close to solar activity. Astrophysics. V. 50, No 2, pp. 187–193.

Bruevich E.A., & Kononovich E.V. Solar and Solar-type Stars Atmosphere's Activity at 11-year and Quasi-biennial Time Scales. arXiv:1102.3976v1 (2011).

Chumak, O.V., & Matveychuk, T.V. 2010, Forecast of the key parameters of the 24-th solar cycle. Research in Astron. and Astroph. V. 10, No 9, pp. 935–941.

Fligge, M., & Solanki, S.K., & Unruh, Y.C., & Frohlich, C., & Wehrli, C. 1998, A model of solar total and spectral irradiance variations. Astronomy & Astrophys. V. 335, pp. 709–718.

Floyd, L., & Newmark, J., & Cook, J., & Herring, L., & McMullin, D. 2005, Solar EUV and UV spectral irradiances and solar indices. Journal of Atmospheric and Solar-Terrestrial Physics. V. 67 pp. 3–15.

Ishkov, V.N. 2009, 1st Workshop on the activity cycles on the Sun and stars, Moscow, 18-10 December, Edited by EAAO, St-Petersburg, pp. 57.

Kleczek, J. 1952, Catalogue de l'activite' des e'ruptions chromosphe'riques. Publ. Inst. Centr. Astron., No. 22. (Czechoslovakia, Prague: Inst. Centr. Astron.)

Krivova, N.A., & Solanki, S.K., & Fligge, M., & Unruh, Y. C. 2003, Reconstruction of solar total and spectral irradiance variations in cycle 23: is solar surface magnetism the cause? Astron. Astrophys. V. 399, L1–L4.

Krivova, N. A., & Solanki, S. K. 2008, Models of solar irradiance variations: current status. Journal of Astrophysics and Astronomy. V. 29, pp. 151—158.

Lukyanova, R., & Mursula, K. 2011, Changed relation between sunspot numbers, solar UV/EUV radiation and TSI during the declining phase of solar cycle 23. Journal of Atmospheric and Solar-Terrestrial Physics. V. 73 pp. 235–240.

National Geophysical Data Center. Solar-Geophysical Data Reports. 54 Years of Space Weather Data. 2009, //www.ngdc.noaa.gov/stp/solar/sgd.html.

National Geophysical Data Center. Solar Data Service. Sun, solar activity and upper atmosphere data. 2012, //www.ngdc.noaa.gov/stp/solar/solardataservices.html.

Viereck, R., & Puga, L., & McMullin, D., & Judge, D., & Weber, M. & Tobiska, K. The MgII index: a proxy for solar EUV. Journal of Geophysical Research, V. 28, No 7, pp. 1343–1346.

Skupin, J., & Noyel, S.,& Wuttke, M.W., & Gottwald, M., & Bovensmann, H., & Weber, M., & Burrows, J. P. 2005, SCIAMACHY solar irradiance observation in the spectral range from 240 to 2380 nm . Advance Space Res. V. 35, pp. 370–375.

Viereck, R.A., Floyd, L.E., Crane, P.C., Woods, T.N., Knapp, B.G., Rottman, G., Weber, M., Puga, L.C. and Deland, M.T. A composite MgII index spanning from 1978 to 2003. Space Weather 2, No. 10, doi:10.1029/2004SW000084.

Vitinsky, Yu.,& Kopezky, M., & Kuklin G., 1986. The sunspot solar activity statistik. M. Nauka



Figure 4: Correlation between monthly averages of solar indices versus $F_{10,7}$ in the cycles 21 - 23. (a) Wolf numbers, (b) $F_{530,3}$, (c) MgII core-to-wing ratio, (d) Flare Index, (e) Counts of flares and (f) TSI.



Figure 5: Correlation coefficients of linear regression $K_{corr}(t)$ for (a) TSI, (b) Flare Index, (c) Counts of flares and (d) MgII UV-index versus W and $F_{10,7}$. $K_{corr}(t)$ was calculated during three-year interval. Solar cycles 21 - 23.